

# Food Allergy Action Plan

## Emergency Care Plan

Place  
Camper's  
Picture  
Here

Name: \_\_\_\_\_ D.O.B.: \_\_\_\_/\_\_\_\_/\_\_\_\_

Allergy to: \_\_\_\_\_

Weight: \_\_\_\_\_ lbs. Asthma:  Yes (higher risk for a severe reaction)  No

Extremely reactive to the following foods: \_\_\_\_\_

### THEREFORE:

- If checked, give epinephrine immediately for ANY symptoms if the allergen was *likely* eaten.
- If checked, give epinephrine immediately if the allergen was *definitely* eaten, even if no symptoms are noted.

### Any SEVERE SYMPTOMS after suspected or known ingestion:

One or more of the following:

- LUNG: Short of breath, wheeze, repetitive cough
- HEART: Pale, blue, faint, weak pulse, dizzy, confused
- THROAT: Tight, hoarse, trouble breathing/swallowing
- MOUTH: Obstructive swelling (tongue and/or lips)
- SKIN: Many hives over body

Or combination of symptoms from different body areas:

- SKIN: Hives, itchy rashes, swelling (e.g., eyes, lips)
- GUT: Vomiting, diarrhea, crampy pain



### 1. INJECT EPINEPHRINE IMMEDIATELY

2. Call 911
3. Begin monitoring (see box below)
4. Give additional medications:\*
  - Antihistamine
  - Inhaler (bronchodilator) if asthma

\*Antihistamines & inhalers/bronchodilators are not to be depended upon to treat a severe reaction (anaphylaxis). USE EPINEPHRINE.

### MILD SYMPTOMS ONLY:

- MOUTH: Itchy mouth
- SKIN: A few hives around mouth/face, mild itch
- GUT: Mild nausea/discomfort



### 1. GIVE ANTIHISTAMINE

2. Stay with camper; alert healthcare professionals and parent
3. If symptoms progress (see above), USE EPINEPHRINE
4. Begin monitoring (see box below)

### Medications/Doses

Epinephrine (brand and dose): \_\_\_\_\_

Antihistamine (brand and dose): \_\_\_\_\_

Other (e.g., inhaler-bronchodilator if asthmatic): \_\_\_\_\_

### Monitoring

**Stay with camper; alert healthcare professionals and parent.** Tell rescue squad epinephrine was given; request an ambulance with epinephrine. Note time when epinephrine was administered. A second dose of epinephrine can be given 5 minutes or more after the first if symptoms persist or recur. For a severe reaction, consider keeping camper lying on back with legs raised. Treat camper even if parents cannot be reached. See back/attached for auto-injection technique.

Parent/Guardian Signature \_\_\_\_\_

Date \_\_\_\_\_


Physician/Healthcare Provider Signature \_\_\_\_\_

Date \_\_\_\_\_


TURN FORM OVER

## EP PEN Auto injector and EP PEN Jr Auto injector Directions

- First, remove the EP PEN Auto injector from the plastic carrying case
- Pull off the blue safety release cap



- Hold orange tip near outer thigh (a ways apply to thigh)



- Swing and firmly push orange tip against outer thigh. Hold on thigh for approximately 10 seconds. Remove the EP PEN Auto injector and massage the area for 10 more seconds



DEY™ and the Dey logo, EpiPen™, EpiPen 2-Pak™, and EpiPen Jr 2-Pak™ are registered trademarks of Dey Pharma, L.P.

## Adrenaclick™ 0.3 mg and Adrenaclick™ 0.15 mg Directions



Remove GREY caps labeled "1" and "2."



Place RED rounded tip against outer thigh, press down hard until needle penetrates. Hold for 10 seconds, then remove.

A food allergy response kit should contain at least two doses of epinephrine, other medications as noted by the camper's physician, and a copy of this Food Allergy Action Plan.

A kit must accompany the camper.

## Contacts

Doctor: \_\_\_\_\_

Phone: (\_\_\_\_) \_\_\_\_ - \_\_\_\_\_

Parent/Guardian: \_\_\_\_\_

Phone: (\_\_\_\_) \_\_\_\_ - \_\_\_\_\_

## Other Emergency Contacts

Name/Relationship: \_\_\_\_\_

Phone: (\_\_\_\_) \_\_\_\_ - \_\_\_\_\_

Name/Relationship: \_\_\_\_\_

Phone: (\_\_\_\_) \_\_\_\_ - \_\_\_\_\_